




# SMALL FLOCK PROGRAM



NOTE: Feed consumptions will vary according to health status, genetics, feeding practices and other management factors.

PURINATURE Consumption Chart	0 to 4 weeks	4 to 8 weeks	8 to 10 weeks	10 to 14 weeks	14 to 18 weeks	18 weeks +
	BROILERS - ROASTERS	2 kg	4 kg	4 kg +		
(With 6054-6055)	6 kg		4 kg +			
TURKEYS	3 kg	5 kg	11 kg			
LAYERS	3 kg		6 kg			

### FEEDING PROGRAM FOR BROILERS/ROASTERS

Chicks should be fed on a free-choice basis and should have fresh feed at all times. After 5 weeks of age, double the feeder space and water capacity.

**NOTE:** Broilers are usually processed at about 2 kg (4.4 lbs) while roasters can be processed at 3.5 kg (8 lbs). Pullets are usually smaller and lighter than cockerels of the same age.

### FEEDING PROGRAM FOR TURKEYS

Provide one tube feeder for every 10-12 turkeys. If trough feeders are used, 20 cm (8 in.) are required per bird. Turkey poults must have easy access to feed during the first week of life. Try using egg cartons, since they make excellent feeders for starting turkeys. Clean fresh water should be available for the birds on a free-choice basis at all times. At 6 weeks of age, increase the floor space up to 0.5 sq. m (5 sq. ft.) per bird and double the feeder space and water capacity.

**NOTE:** Turkeys will achieve a weight of about 5 kg (11 lbs) at about 14 weeks but toms can be grown to more than 13 kg (29 lbs) in 25 weeks. Hens are generally smaller at the same age.

### FEEDING PROGRAM FOR EGG LAYING BIRDS

Birds should be fed on a free-choice basis fed at all times and fresh, room temperature water should always be available. After 20 weeks, feeding and watering space should be increased to one tube feeder for every 10-15 hens or 10 cm (4 in) feeder space per hen. There should be one nest for every 5-7 hens. Supplemental light should be provided. Ensure that the hens receive 14-17 hours of continuous light per day to stay in egg production. If the lights are turned on at 6 am, most hens will lay their eggs by 11 am. Supplemental heat should be provided when the temperature is less than 15°C (60°F) to maintain egg production.

GUARANTEED ANALYSIS						
	Protein	Fat	Fibre	Na	Ca	P
6032	15%	2.0%	6.0%	0.16%	0.85%	0.70%
6038	20%	4.0%	4.5%	0.18%	1.00%	0.75%
6039	20%	4.0%	4.5%	0.18%	1.00%	0.75%
6054	18%	3.0%	5.0%	0.16%	0.90%	0.70%
6498/6498 "L"	17%	1.5%	4.5%	0.16%	3.80%	0.60%
8016	24%	3.0%	6.0%	0.15%	1.30%	0.90%
8036	20%	6.0%	6.0%	0.15%	1.10%	0.85%
8070	25%	2.0%	6.0%	0.15%	1.30%	0.90%
8071	20%	2.0%	6.0%	0.16%	1.00%	0.80%

[multipurina.ca](http://multipurina.ca)

For more information, contact your local Purina Dealer: 1-800-567-CHOW(2469), [purina\\_retail@cargill.com](mailto:purina_retail@cargill.com) or visit our website.